

## SCA54

### Decreased Cerebral Tissue Oxygen Saturation during Aortic Surgery Increases Risk of Post-Operative Complications

Fischer G; Lin H; DiLuozzo G; Griep R; Reich D  
*Mount Sinai School of Medicine, New York, New York, USA*

**Introduction:** Currently only limited information exists exploring the relationship between brain oxygenation levels during aortic surgery and post-operative outcomes (1). The FORE-SIGHT® (CAS Medical Systems, Branford CT USA) cerebral oximeter measures absolute cerebral tissue oxygen saturation (SctO<sub>2</sub>) values. We hypothesized that a relationship exists between decreased intraoperative SctO<sub>2</sub> values and post-operative complications following aortic surgery.

**Methods:** With IRB approval and informed consent, patients undergoing elective thoracic aortic surgery with deep hypothermic circulatory arrest (DHCA) were monitored intraoperatively using the FORE-SIGHT. Two sensors were placed on the subject's forehead bilaterally with SctO<sub>2</sub> values recorded every 2 seconds starting after induction of anesthesia until the end of surgery. SctO<sub>2</sub> minutes and the area above the curve (AAC) spent beneath the absolute threshold limits of 50, 55, 58, 60, & 65% were computed for both left & right sensors. Complications were categorized as "major" (death, stroke, depressed LV function, respiratory failure, sepsis, delirium, renal failure, GI complications, or severe volume overload) and "minor" (atrial fibrillation, minor volume overload, phlebitis, or none). Post-operative complications, Extubation time, ICU length of stay, and Hospital length of stay (HLOS) data were collected and compared to SctO<sub>2</sub> data and DHCA time.

**Results:** Demographics: 30 subjects; Gender 22M/8F; Race 25W/1AA/3HS/1IN; Post-induction SctO<sub>2</sub> was 70.6% (SD 5.1). SctO<sub>2</sub><60% (minutes & AUC), SctO<sub>2</sub><58% (minutes), and DHCA minutes were significantly associated with Extubation time and ICU LOS ( $p<0.05$  two-tailed, Spearman). Major complications significantly correlated ( $p<0.05$  two-tailed, Spearman) to Extubation time (median: 3 days vs 1 day for subjects with major complication) and HLOS (median: 13 days vs 7.5 days). Logistic regression odds ratio analysis showed that for every increment of 30 minutes below a SctO<sub>2</sub> threshold, the risk for major complications increased 2.1 times for SctO<sub>2</sub><60% and 3.7 times for SctO<sub>2</sub><55% (Table 1). Similar results were found for AAC of SctO<sub>2</sub> thresholds with 50 minute-% increments. Also for every 5 minutes of DHCA, the risk of major complications increased 1.6 times.

**Discussion:** Despite the low number of subjects enrolled in this study, decreased SctO<sub>2</sub> values and prolonged DHCA times were found to be associated with major complications, prolonged extubation times, and ICU/Hospital LOS. This study suggests that prolonged intraoperative periods of time with SctO<sub>2</sub><60% correlate significantly with an increased risk of having major complications and an increased LOS.

This study was partially supported by NIH grant NS045488.

1) Reich et al, Eur J Cardiothorac Surg. 2004 Mar;25(3):401-6.

<b>Table 1 Brain hemisphere with lower SctO2</b>	<b>Minor or No Complication (n=14)</b>	<b>Major Complication (n=16)</b>	<b>Odds ratio unit</b>	<b>Odds ratio* (95% CI)</b>	<b>p-value of odds ratio*</b>
	Median (IQR)	Median (IQR)			
<b>SctO2 &lt; 60% (minutes)</b>	2.3 (0 - 25.4)	59.3 (3.9-128.5)	30 minutes	2.11 (1.18, 5.16)	<0.01
<b>SctO2 &lt; 58% (minutes)</b>	1.6 (0 - 14.3)	34.9 (0.8 - 94.7)		2.58 (1.18, 8.32)	<0.01
<b>SctO2 &lt; 55% (minutes)</b>	0.45 (0 - 9.5)	10.9 (0 - 54.8)		3.71 (1.04, 33.78)	0.03
<b>SctO2 &lt; 50% (minutes)</b>	0 (0 - 0.21)	0 (0 - 13.7)		219.20 (1.03, >999)	0.04
<b>SctO2 &lt; 60% (AUC)</b>	8.1 (0 - 115.1)	200.8 (5.4- 648.8)	50 minutes- %	1.61 (1.05, 3.00)	0.01
<b>SctO2 &lt; 58% (AUC)</b>	4.1 (0 - 72.1)	101.7 (1.7- 420.6)		2.01 (1.04, 5.81)	0.02
<b>SctO2 &lt; 55% (AUC)</b>	0.55 (0 - 18.8)	15.9 (0 - 195.2)		3.22 (1.01, 36.60)	0.04
<b>SctO2 &lt; 50% (AUC)</b>	0 (0 - 0.57)	0 (0 - 41.9)		>999 (0, >999)	0.02
<b>DHCA duration (minutes)</b>	21.0 (18.8 - 23.8)	26.5 ( 20.0 - 33.75)	5 minutes	1.92 (1.05, 4.13)	0.03